

Dear Author,

Here are the final proofs of your article. Please check the proofs carefully.

Please note that at this stage you should only be checking for errors introduced during the production process. Please pay particular attention to the following when checking the proof:

- Author names. Check that each author name is spelled correctly, and that names appear in the correct order of first name followed by family name. This will ensure that the names will be indexed correctly (for example if the author's name is 'Patel, J. ', she will be cited as ' Jane Patel ').
- Affiliations. Check that all authors are cited with the correct affiliations, that the author who will receive correspondence has been identified with an asterisk (\*), and that all equal contributors have been identified with a well sign (#).
- Ensure that the main text is complete.
- Check that figures, tables and their legends are included and in the correct order.
- Look to see that queries that were raised during copy-editing or typesetting have been resolved.
- Confirm that all web links are correct and working.
- Ensure that special characters and equations are displaying correctly.
- Check that additional or supplementary files can be opened and are correct.

Changes in scientific content cannot be made at this stage unless the request has already been approved. This includes changes to title or authorship, new results, or corrected values.

## How to return your corrections

Returning your corrections via email:

- Annotate the proof PDF with your corrections.
- Remember to include the journal title, manuscript number, and your name when sending your response via email.

After you have submitted your corrections, you will receive email notification from our production team that your article has been published in the final version. All changes at this stage are final. We will not be able to make any further changes after publication.

Kind regards,



Case Report

Creative Commons 4.0

# A special discovery of pancreatic adenocarcinoma by Sister Mary-Joseph's Nodule: what is the prognosis?

Mohamed Ali Chaouch<sup>a</sup>\*, Asma Chaouch<sup>a</sup>, Karim Nacef<sup>a</sup>, Moez Boudokhane<sup>a</sup>

#### **Abstract**

Cutaneous metastases of visceral tumours are uncommon and might have umbilical and/or extra umbilical locations. The umbilical location, also named «Nodule of Sister Mary-Joseph» among 7 to 9% is secondary to a pancreatic tumour. In this article, we reported a case of a 50-years-old man presenting pancreatic cancer discovered by umbilical nodules and weight loss noticed 3 months earlier treated by palliative chemotherapy. We report this case of a rare presentation of cutaneous metastases of pancreatic disease to insist on its bad prognosis level.

Keywords: Cutaneous metastasis; pancreatic cancer; chemotherapy; nodule of Sister Mary Joseph

#### INTRODUCTION

Cutaneous metastases of visceral tumours are uncommon. It can be umbilical and or extra umbilical. The umbilical location is known as the "Nodule of Sister Mary-Joseph" (NSMJ). It is described for the first time in 1949 by Sir Hamilton Bailey [1] and since about 600 cases of NSMJ have been published which only 7 to 9% are secondary to a pancreatic tumour [2]. We report this case of a rare presentation of cutaneous metastases of pancreatic disease to insist on its bad prognosis level.

## **CASE REPORT**

A 50-year-old men, with a familial history of an idiopathic mega-oesophagus in the father and adenocarcinoma of the prostate in the paternal uncle, consult for umbilical nodules and a weight loss for 3 months. Physical examination reveals an irregular, indurated, umbilical nodules measuring 2 x 1.5 cm (Figure 1). Biological exams show anaemia at 9 g / dl. Abdominal ultrasound shows an umbilical hypoechoic heterogeneous non-vascularized mass.

An abdominal CT scan revealed a hypodense tissue mass

at the body and tail of the pancreas measuring 59 x 48 x 44 cm with fuzzy limits and little enhancement after injection with heterogeneous necrotic appearance (Figure 2). This invades the splenic vein. Hypodense epiploic masses of 26 mm per pancreatic in contact with the gastric wall and 11.4 mm peri-umbilical suggest nodules of carcinomatosis. The patient had many biopsies of this umbilical lesion. Anatomopathological examination with an immunohistochemical study concluded with cutaneous metastasis of pancreatic adenocarcinoma. The diagnosis of a tumour of the body and the tail of the pancreas with umbilical and peritoneal metastases was



Figure 1. An umbilical nodule of hard consistency, with nipple surface, ulcerated in places and not painful.

E-mail: Docmedalichaouch@gmail.com

Received: 24 December 2019 Accepted: 12 February 2020

<sup>&</sup>lt;sup>a</sup>Department of General Surgery, Tahar Sfar Hospital, Mahdia, Tunisia.

<sup>\*</sup>Corresponding author: Dr. Mohamed Ali Chaouch

<sup>🔟</sup> ing address: Department of general surgery, Tahar Sfar 🞝 ital, Mahdia, Tunisia.




Figure 2. An abdominal CT-scan showing a necrotic pancreatic tumor (white arrow).

retained. The patient had palliative chemotherapy with Cisplatin. The patient was dead after six months.

## **DISCUSSION**

Sister Mary-Joseph (1856-1939) was an assistant nurse to the surgeon William J. Mayo, son of William Worrall Mayo, founder of the Mayo Clinic in Rochester, was the first to notice the association of an umbilical nodule with an abdominal tumour during the examination of patients undergoing laparotomy [3]. These nodules are present in 1 to 3% of patients with abdominal tumour [4]. They represent 30% of umbilical tumours and they are witnesses of adenocarcinoma in 90% of cases [5]. But they can also be associated with squamous cell carcinoma, melanoma or sarcoma. The gastrointestinal tract is the most common origin of the primary tumour followed by gynaecological localization [6]. The most-reported digestive origins in descending order are stomach (26%), colon (10%) and pancreas (7%)  $^{[6]}$ . Primary tumor cannot be found in 15% of cases [7,8]. The pathophysiology remains unclear and several hypotheses have been proposed. Direct extension of peritoneal lesions remains the most common mechanism. Hematogenous dissemination or lymphatic extension along with ligaments of embryonic or iatrogenic origin during laparoscopy is reported [9]. It concerns essentially adults with equal sex ratio [2]. It is known that pancreatic tumor remains asymptomatic for a long time. Then, they are usually discovered at an advanced stage [2, 11]. The umbilical cutaneous metastasis

is seen more frequently than other cutaneous localizations. Patient survival only a few months <sup>[2]</sup>. Admittedly, the NSMJ is a sign of neoplastic progression or recurrence remains a sign prognosis pejorative even suggesting the therapeutic abstention. However, some studies <sup>[9,12]</sup>, confirming our results, have led to the conclusion that early diagnosis with combined surgery-chemotherapy seems to provide a better survival response for pancreatic tumors discovered as a result of NSMJ. Doctors should know this rare and characteristic umbilical nodule correlated with a bad prognosis. Their clinical characteristics do not generally make it possible to differentiate them from the primitive lesions with the naked eye. But their superficial seat makes them accessible to a biopsy which can facilitate the diagnosis.

## **DECLARATIONS**

## Acknowledgements

None

## Authors' contributions

Dr Mohamed Ali Chaouch: Participate in surgical treatment and writing the article.

Dr Asma Chaouch: Follow the patient in the outcome point Dr Karim Nacef and Dr Moez Boudokhane: Checked the article and supervised the patient's care.

## Availability of Data and Materials

Not applicable

#### Financial support and sponsorship

None

#### Conflicts of interest

The authors have no conflict of interest to disclose

## Ethical approval and consent to participate

Not applicable

#### Consent for publication

Not applicable

## REFERENCES

- Abu-Hilal, M., & Newman, J. S. (2009). Sister Mary Joseph and her nodule: historical and clinical perspective. The American journal of the medical sciences, 337(4), 271-273.
- Yendluri, V., Centeno, B., & Springett, G.M. (2007). Pancreatic cancer presenting as a Sister Mary Joseph's nodule: case report and update of the literature. Pancreas, 34(1), 161-164
- 3. J., Jacques, L., Mesturoux, C., & Vong, et al. (2014). Un nodule ombilical. La Revue de Médecine Interne.
- 4. Papa Souleymane Touré, Tall, C. T., Pauline Dioussé, Adama Berthé, & Ka, M. M.. (2015). Nodule de sœur mariejosèphe révélateur de carcinomes digestif et ovarien: à propos de 4 cas. Pan African Medical Journal, 22.
- 5. Khalfallah, M., Chaker, Y., & Dziri, C.. (2011). Sister mary joseph's nodule showing adenocarcinoma of pancreas. La

- Tunisie médicale, 89(10), 790-791.
- Gharaba, S., Elfadil, K., Samlani, Z., Difaa, A., Krati, K., & Elmahfoudi, A., et al. (2011). Nodule de s?ur-marie-joseph: quelles implications diagnostiques et thérapeutiques?
  à propos de deux cas. Journal Africain d'Hépato-Gastroentérologie, 5(4), 276-279.
- 7. Chalya, P. L., Mabula, J. B., & Peter F Rambau.... (2013). Sister mary joseph's nodule at a university teaching hospital in northwestern tanzania: a retrospective review of 34 cases. World Journal of Surgical Oncology, 11(1), 151.
- 8. Gilbert, J. D., & Byard, R. W. (2016). Sister mary joseph's nodule: a very useful indicator of significant internal pathology. Forensic Science, Medicine, and Pathology, 12(1), 101-103.
- 9. Ozaki, N., Takamori, H., & Baba, H.. (2011). Sister mary joseph's nodule derived from pancreatic cancer. Journal of Hepato-Biliary-Pancreatic Sciences, 18(1), 119-121.
- Zhou, Wang, Gao, Bu, Zhang, & Wang. (2014). Cutaneous metastasis from pancreatic cancer: a case report and systematic review of the literature. Oncology Letters.
- 11. Lau, Melvin K., Davila, Jessica A., & Shaib, Yasser H. . Incidence and survival of pancreatic head and body and tail cancers. Pancreas, 39(4), 458-462.
- Bai, X. L., Zhang, Q., Masood, W., Masood, N., Tang, Y., Cao,
  C. H. & Liang, T. B. (2012). Sister Mary Joseph's nodule
  as a first sign of pancreatic cancer. World Journal of
  Gastroenterology: WJG, 18(45), 6686.



## Author Query Form

## Dear Author,

During the copy-editing of your paper, the following queries arose. Please refer to the query reference call out numbers in the page proofs and respond to each by marking the necessary comments using the PDF annotation tools. Please remember illegible or unclear comments and corrections may delay publication.

Many thanks for your assistance.

Query Reference	Query	Remark
Q1	Author: Please confirm that given names and surnames/family names have been identified correctly.	
Q2	Affiliations: Please check if the affiliations are presented correctly.	
Q3	Corresponding author: Please check if the information are presented correctly.	
Q4	Please check if the reference is correct? There is no 10th reference in the main text?	
Q5	Please check if the chart annotation is correct?	